

Search for charged Higgs bosons in decays of top quarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,²⁴ J. Adelman,¹⁴ T. Akimoto,⁵⁶ B. Álvarez González^{q,12} S. Amerio^{w,44} D. Amidei,³⁵ A. Anastassov,³⁹ A. Annovi,²⁰ J. Antos,¹⁵ G. Apollinari,¹⁸ A. Apresyan,⁴⁹ T. Arisawa,⁵⁸ A. Artikov,¹⁶ W. Ashmanskas,¹⁸ A. Attal,⁴ A. Aurisano,⁵⁴ F. Azfar,⁴³ P. Azzurri^{z,47} W. Badgett,¹⁸ A. Barbaro-Galtieri,²⁹ V.E. Barnes,⁴⁹ B.A. Barnett,²⁶ V. Bartsch,³¹ G. Bauer,³³ P.-H. Beauchemin,³⁴ F. Bedeschi,⁴⁷ D. Beecher,³¹ S. Behari,²⁶ G. Bellettini^{x,47} J. Bellinger,⁶⁰ D. Benjamin,¹⁷ A. Beretvas,¹⁸ J. Beringer,²⁹ A. Bhatti,⁵¹ M. Binkley,¹⁸ D. Bisello^{w,44} I. Bizjak^{cc,31} R.E. Blair,² C. Blocker,⁷ B. Blumenfeld,²⁶ A. Bocci,¹⁷ A. Bodek,⁵⁰ V. Boisvert,⁵⁰ G. Bolla,⁴⁹ D. Bortoletto,⁴⁹ J. Boudreau,⁴⁸ A. Boveia,¹¹ B. Brau^{a,11} A. Bridgeman,²⁵ L. Brigliadori,⁴⁴ C. Bromberg,³⁶ E. Brubaker,¹⁴ J. Budagov,¹⁶ H.S. Budd,⁵⁰ S. Budd,²⁵ S. Burke,¹⁸ K. Burkett,¹⁸ G. Busetto^{w,44} P. Bussey,²² A. Buzatu,³⁴ K. L. Byrum,² S. Cabrera^{s,17} C. Calancha,³² M. Campanelli,³⁶ M. Campbell,³⁵ F. Canelli^{14,18} A. Canepa,⁴⁶ B. Carls,²⁵ D. Carlsmith,⁶⁰ R. Carosi,⁴⁷ S. Carrillo^{l,19} S. Carron,³⁴ B. Casal,¹² M. Casarsa,¹⁸ A. Castro^{v,6} P. Catastini^{y,47} D. Cauz^{bb,55} V. Cavaliere^{y,47} M. Cavalli-Sforza,⁴ A. Cerri,²⁹ L. Cerrito^{m,31} S.H. Chang,²⁸ Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁷ G. Chlachidze,¹⁸ F. Chlebana,¹⁸ K. Cho,²⁸ D. Chokheli,¹⁶ J.P. Chou,²³ G. Choudalakis,³³ S.H. Chuang,⁵³ K. Chung,¹³ W.H. Chung,⁶⁰ Y.S. Chung,⁵⁰ T. Chwalek,²⁷ C.I. Ciobanu,⁴⁵ M.A. Ciocci^{y,47} A. Clark,²¹ D. Clark,⁷ G. Compostella,⁴⁴ M.E. Convery,¹⁸ J. Conway,⁸ M. Cordelli,²⁰ G. Cortiana^{w,44} C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{x,47} C. Cuenca Almenar^{s,8} J. Cuevas^{q,12} R. Culbertson,¹⁸ J.C. Cully,³⁵ D. Dagenhart,¹⁸ M. Datta,¹⁸ T. Davies,²² P. de Barbaro,⁵⁰ S. De Cecco,⁵² A. Deisher,²⁹ G. De Lorenzo,⁴ M. Dell'Orso^{x,47} C. Deluca,⁴ L. Demortier,⁵¹ J. Deng,¹⁷ M. Deninno,⁶ P.F. Derwent,¹⁸ G.P. di Giovanni,⁴⁵ C. Dionisi^{aa,52} B. Di Ruzza^{bb,55} J.R. Dittmann,⁵ M. D'Onofrio,⁴ S. Donati^{x,47} P. Dong,⁹ J. Donini,⁴⁴ T. Dorigo,⁴⁴ S. Dube,⁵³ J. Efron,⁴⁰ A. Elagin,⁵⁴ R. Erbacher,⁸ D. Errede,²⁵ S. Errede,²⁵ R. Eusebi,¹⁸ H.C. Fang,²⁹ S. Farrington,⁴³ W.T. Fedorko,¹⁴ R.G. Feild,⁶¹ M. Feindt,²⁷ J.P. Fernandez,³² C. Ferrazza^{z,47} R. Field,¹⁹ G. Flanagan,⁴⁹ R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²³ J.C. Freeman,¹⁸ I. Furic,¹⁹ M. Gallinaro,⁵² J. Galyardt,¹³ F. Garbersen,¹¹ J.E. Garcia,²¹ A.F. Garfinkel,⁴⁹ K. Genser,¹⁸ H. Gerberich,²⁵ D. Gerdes,³⁵ A. Gessler,²⁷ S. Giagu^{aa,52} V. Giakoumopoulou,³ P. Giannetti,⁴⁷ K. Gibson,⁴⁸ J.L. Gimmell,⁵⁰ C.M. Ginsburg,¹⁸ N. Giokaris,³ M. Giordani^{bb,55} P. Giromini,²⁰ M. Giunta^{x,47} G. Giurgiu,²⁶ V. Glagolev,¹⁶ D. Glenzinski,¹⁸ M. Gold,³⁸ N. Goldschmidt,¹⁹ A. Golossanov,¹⁸ G. Gomez,¹² G. Gomez-Ceballos,³³ M. Goncharov,³³ O. González,³² I. Gorelov,³⁸ A.T. Goshaw,¹⁷ K. Goulianos,⁵¹ A. Gresele^{w,44} S. Grinstein,²³ C. Grosso-Pilcher,¹⁴ R.C. Group,¹⁸ U. Grundler,²⁵ J. Guimaraes da Costa,²³ Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S.R. Hahn,¹⁸ E. Halkiadakis,⁵³ B.-Y. Han,⁵⁰ J.Y. Han,⁵⁰ F. Happacher,²⁰ K. Hara,⁵⁶ D. Hare,⁵³ M. Hare,⁵⁷ S. Harper,⁴³ R.F. Harr,⁵⁹ R.M. Harris,¹⁸ M. Hartz,⁴⁸ K. Hatakeyama,⁵¹ C. Hays,⁴³ M. Heck,²⁷ A. Heijboer,⁴⁶ J. Heinrich,⁴⁶ C. Henderson,³³ M. Herndon,⁶⁰ J. Heuser,²⁷ S. Hewamanage,⁵ D. Hidas,¹⁷ C.S. Hill^{c,11} D. Hirschbuehl,²⁷ A. Hocker,¹⁸ S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,²⁹ B.T. Huffman,⁴³ R.E. Hughes,⁴⁰ U. Husemann,⁶¹ M. Hussein,³⁶ J. Huston,³⁶ J. Incandela,¹¹ G. Introzzi,⁴⁷ M. Iori^{aa,52} A. Ivanov,⁸ E. James,¹⁸ D. Jang,¹³ B. Jayatilaka,¹⁷ E.J. Jeon,²⁸ M.K. Jha,⁶ S. Jindariani,¹⁸ W. Johnson,⁸ M. Jones,⁴⁹ K.K. Joo,²⁸ S.Y. Jun,¹³ J.E. Jung,²⁸ T.R. Junk,¹⁸ T. Kamon,⁵⁴ D. Kar,¹⁹ P.E. Karchin,⁵⁹ Y. Kato,⁴² R. Kephart,¹⁸ J. Keung,⁴⁶ V. Khotilovich,⁵⁴ B. Kilminster,¹⁸ D.H. Kim,²⁸ H.S. Kim,²⁸ H.W. Kim,²⁸ J.E. Kim,²⁸ M.J. Kim,²⁰ S.B. Kim,²⁸ S.H. Kim,⁵⁶ Y.K. Kim,¹⁴ N. Kimura,⁵⁶ L. Kirsch,⁷ S. Klimentenko,¹⁹ B. Knuteson,³³ B.R. Ko,¹⁷ K. Kondo,⁵⁸ D.J. Kong,²⁸ J. Konigsberg,¹⁹ A. Korytov,¹⁹ A.V. Kotwal,¹⁷ M. Kreps,²⁷ J. Kroll,⁴⁶ D. Krop,¹⁴ N. Krumnack,⁵ M. Kruse,¹⁷ V. Krutelyov,¹¹ T. Kubo,⁵⁶ T. Kuhr,²⁷ N.P. Kulkarni,⁵⁹ M. Kurata,⁵⁶ S. Kwang,¹⁴ A.T. Laasanen,⁴⁹ S. Lami,⁴⁷ S. Lammel,¹⁸ M. Lancaster,³¹ R.L. Lander,⁸ K. Lannon^{p,40} A. Lath,⁵³ G. Latino^{y,47} I. Lazzizzera^{w,44} T. LeCompte,² E. Lee,⁵⁴ H.S. Lee,¹⁴ S.W. Lee^{r,54} S. Leone,⁴⁷ J.D. Lewis,¹⁸ C.-S. Lin,²⁹ J. Linacre,⁴³ M. Lindgren,¹⁸ E. Lipeles,⁴⁶ A. Lister,⁸ D.O. Litvintsev,¹⁸ C. Liu,⁴⁸ T. Liu,¹⁸ N.S. Lockyer,⁴⁶ A. Loginov,⁶¹ M. Loretz^{w,44} L. Lovas,¹⁵ D. Lucchesi^{w,44} C. Luci^{aa,52} J. Lueck,²⁷ P. Lujan,²⁹ P. Lukens,¹⁸ G. Lungu,⁵¹ L. Lyons,⁴³ J. Lys,²⁹ R. Lysak,¹⁵ D. MacQueen,³⁴ R. Madrak,¹⁸ K. Maeshima,¹⁸ K. Makhoul,³³ T. Maki,²⁴ P. Maksimovic,²⁶ S. Malde,⁴³ S. Malik,³¹ G. Manca^{e,30} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁹ C. Marino,²⁷ C.P. Marino,²⁵ A. Martin,⁶¹ V. Martin^{k,22} M. Martínez,⁴ R. Martínez-Ballarín,³² T. Maruyama,⁵⁶ P. Mastrandrea,⁵² T. Masubuchi,⁵⁶ M. Mathis,²⁶ M.E. Mattson,⁵⁹ P. Mazzanti,⁶ K.S. McFarland,⁵⁰ P. McIntyre,⁵⁴ R. McNulty^{j,30} A. Mehta,³⁰ P. Mehtala,²⁴ A. Menzione,⁴⁷ P. Merkel,⁴⁹ C. Mesropian,⁵¹ T. Miao,¹⁸ N. Miladinovic,⁷ R. Miller,³⁶ C. Mills,²³ M. Milnik,²⁷ A. Mitra,¹ G. Mitselmakher,¹⁹ H. Miyake,⁵⁶ N. Moggi,⁶ C.S. Moon,²⁸ R. Moore,¹⁸ M.J. Morello^{x,47}

J. Morlock,²⁷ P. Movilla Fernandez,¹⁸ J. Mülmenstädt,²⁹ A. Mukherjee,¹⁸ Th. Muller,²⁷ R. Mumford,²⁶ P. Murat,¹⁸ M. Mussini^v,⁶ J. Nachtman,¹⁸ Y. Nagai,⁵⁶ A. Nagano,⁵⁶ J. Naganoma,⁵⁶ K. Nakamura,⁵⁶ I. Nakano,⁴¹ A. Napier,⁵⁷ V. Necula,¹⁷ J. Nett,⁶⁰ C. Neu^t,⁴⁶ M.S. Neubauer,²⁵ S. Neubauer,²⁷ J. Nielsen^g,²⁹ L. Nodulman,² M. Norman,¹⁰ O. Norniella,²⁵ E. Nurse,³¹ L. Oakes,⁴³ S.H. Oh,¹⁷ Y.D. Oh,²⁸ I. Oksuzian,¹⁹ T. Okusawa,⁴² R. Orava,²⁴ K. Osterberg,²⁴ S. Pagan Griso^w,⁴⁴ E. Palencia,¹⁸ V. Papadimitriou,¹⁸ A. Papaikonomou,²⁷ A.A. Paramonov,¹⁴ B. Parks,⁴⁰ S. Pashapour,³⁴ J. Patrick,¹⁸ G. Pauletta^{bb},⁵⁵ M. Paulini,¹³ C. Paus,³³ T. Peiffer,²⁷ D.E. Pellett,⁸ A. Penzo,⁵⁵ T.J. Phillips,¹⁷ G. Piacentino,⁴⁷ E. Pianori,⁴⁶ L. Pinera,¹⁹ K. Pitts,²⁵ C. Plager,⁹ L. Pondrom,⁶⁰ O. Poukhov^{*},¹⁶ N. Pounder,⁴³ F. Prakhoshyn,¹⁶ A. Pronko,¹⁸ J. Proudfoot,² F. Ptohosⁱ,¹⁸ E. Pueschel,¹³ G. Punzi^x,⁴⁷ J. Pursley,⁶⁰ J. Rademacker^c,⁴³ A. Rahaman,⁴⁸ V. Ramakrishnan,⁶⁰ N. Ranjan,⁴⁹ I. Redondo,³² P. Renton,⁴³ M. Renz,²⁷ M. Rescigno,⁵² S. Richter,²⁷ F. Rimondi^v,⁶ L. Ristori,⁴⁷ A. Robson,²² T. Rodrigo,¹² T. Rodriguez,⁴⁶ E. Rogers,²⁵ S. Rolli,⁵⁷ R. Roser,¹⁸ M. Rossi,⁵⁵ R. Rossin,¹¹ P. Roy,³⁴ A. Ruiz,¹² J. Russ,¹³ V. Rusu,¹⁸ H. Saarikko,²⁴ A. Safonov,⁵⁴ W.K. Sakumoto,⁵⁰ O. Saltó,⁴ L. Santi^{bb},⁵⁵ S. Sarkar^{aa},⁵² L. Sartori,⁴⁷ K. Sato,¹⁸ A. Savoy-Navarro,⁴⁵ P. Schlabach,¹⁸ A. Schmidt,²⁷ E.E. Schmidt,¹⁸ M.A. Schmidt,¹⁴ M.P. Schmidt^{*},⁶¹ M. Schmitt,³⁹ T. Schwarz,⁸ L. Scodellaro,¹² A. Scribano^y,⁴⁷ F. Scuri,⁴⁷ A. Sedov,⁴⁹ S. Seidel,³⁸ Y. Seiya,⁴² A. Semenov,¹⁶ L. Sexton-Kennedy,¹⁸ F. Sforza,⁴⁷ A. Sfyrila,²⁵ S.Z. Shalhout,⁵⁹ T. Shears,³⁰ P.F. Shepard,⁴⁸ M. Shimojima^o,⁵⁶ S. Shiraishi,¹⁴ M. Shochet,¹⁴ Y. Shon,⁶⁰ I. Shreyber,³⁷ A. Sidoti,⁴⁷ P. Sinervo,³⁴ A. Sisakyan,¹⁶ A.J. Slaughter,¹⁸ J. Slaunwhite,⁴⁰ K. Sliwa,⁵⁷ J.R. Smith,⁸ F.D. Snider,¹⁸ R. Snihur,³⁴ A. Soha,⁸ S. Somalwar,⁵³ V. Sorin,³⁶ J. Spalding,¹⁸ T. Spreitzer,³⁴ P. Squillacioti^y,⁴⁷ M. Stanitzki,⁶¹ R. St. Denis,²² B. Stelzer,³⁴ O. Stelzer-Chilton,³⁴ D. Stentz,³⁹ J. Strologas,³⁸ G.L. Strycker,³⁵ D. Stuart,¹¹ J.S. Suh,²⁸ A. Sukhanov,¹⁹ I. Suslov,¹⁶ T. Suzuki,⁵⁶ A. Taffard^f,²⁵ R. Takashima,⁴¹ Y. Takeuchi,⁵⁶ R. Tanaka,⁴¹ M. Tecchio,³⁵ P.K. Teng,¹ K. Terashi,⁵¹ J. Thom^h,¹⁸ A.S. Thompson,²² G.A. Thompson,²⁵ E. Thomson,⁴⁶ P. Tipton,⁶¹ P. Tito-Guzmán,³² S. Tkaczyk,¹⁸ D. Toback,⁵⁴ S. Tokar,¹⁵ K. Tollefson,³⁶ T. Tomura,⁵⁶ D. Tonelli,¹⁸ S. Torre,²⁰ D. Torretta,¹⁸ P. Totaro^{bb},⁵⁵ S. Tourneur,⁴⁵ M. Trovato,⁴⁷ S.-Y. Tsai,¹ Y. Tu,⁴⁶ N. Turini^y,⁴⁷ F. Ukegawa,⁵⁶ S. Vallecorsa,²¹ N. van Remortel^b,²⁴ A. Varganov,³⁵ E. Vataga^z,⁴⁷ F. Vázquez^l,¹⁹ G. Velev,¹⁸ C. Vellidis,³ M. Vidal,³² R. Vidal,¹⁸ I. Vila,¹² R. Vilar,¹² T. Vine,³¹ M. Vogel,³⁸ I. Volobouev^r,²⁹ G. Volpi^x,⁴⁷ P. Wagner,⁴⁶ R.G. Wagner,² R.L. Wagner,¹⁸ W. Wagner^u,²⁷ J. Wagner-Kuhr,²⁷ T. Wakisaka,⁴² R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³⁴ D. Waters,³¹ M. Weinberger,⁵⁴ J. Weinelt,²⁷ W.C. Wester III,¹⁸ B. Whitehouse,⁵⁷ D. Whiteson^f,⁴⁶ A.B. Wicklund,² E. Wicklund,¹⁸ S. Wilbur,¹⁴ G. Williams,³⁴ H.H. Williams,⁴⁶ P. Wilson,¹⁸ B.L. Winer,⁴⁰ P. Wittich^h,¹⁸ S. Wolbers,¹⁸ C. Wolfe,¹⁴ T. Wright,³⁵ X. Wu,²¹ F. Würthwein,¹⁰ S. Xie,³³ A. Yagil,¹⁰ K. Yamamoto,⁴² J. Yamaoka,¹⁷ U.K. Yangⁿ,¹⁴ Y.C. Yang,²⁸ W.M. Yao,²⁹ G.P. Yeh,¹⁸ J. Yoh,¹⁸ K. Yorita,⁵⁸ T. Yoshida,⁴² G.B. Yu,⁵⁰ I. Yu,²⁸ S.S. Yu,¹⁸ J.C. Yun,¹⁸ L. Zanello^{aa},⁵² A. Zanetti,⁵⁵ X. Zhang,²⁵ Y. Zheng^d,⁹ and S. Zucchelli^v,⁶

(CDF Collaboration[†])

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^vUniversity of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254*

⁸*University of California, Davis, Davis, California 95616*

⁹*University of California, Los Angeles, Los Angeles, California 90024*

¹⁰*University of California, San Diego, La Jolla, California 92093*

¹¹*University of California, Santa Barbara, Santa Barbara, California 93106*

¹²*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹³*Carnegie Mellon University, Pittsburgh, PA 15213*

¹⁴*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

¹⁵*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁶*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁷*Duke University, Durham, North Carolina 27708*

¹⁸*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

¹⁹*University of Florida, Gainesville, Florida 32611*

²⁰*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

²¹*University of Geneva, CH-1211 Geneva 4, Switzerland*

²²*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²³*Harvard University, Cambridge, Massachusetts 02138*

- ²⁴Division of High Energy Physics, Department of Physics,
University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland
- ²⁵University of Illinois, Urbana, Illinois 61801
- ²⁶The Johns Hopkins University, Baltimore, Maryland 21218
- ²⁷Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
- ²⁸Center for High Energy Physics: Kyungpook National University,
Daegu 702-701, Korea; Seoul National University, Seoul 151-742,
Korea; Sungkyunkwan University, Suwon 440-746,
Korea; Korea Institute of Science and Technology Information, Daejeon,
305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea
- ²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³¹University College London, London WC1E 6BT, United Kingdom
- ³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- ³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
- ³⁴Institute of Particle Physics: McGill University, Montréal, Québec,
Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,
Canada V5A 1S6; University of Toronto, Toronto, Ontario,
Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³⁵University of Michigan, Ann Arbor, Michigan 48109
- ³⁶Michigan State University, East Lansing, Michigan 48824
- ³⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁸University of New Mexico, Albuquerque, New Mexico 87131
- ³⁹Northwestern University, Evanston, Illinois 60208
- ⁴⁰The Ohio State University, Columbus, Ohio 43210
- ⁴¹Okayama University, Okayama 700-8530, Japan
- ⁴²Osaka City University, Osaka 588, Japan
- ⁴³University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴⁴Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^wUniversity of Padova, I-35131 Padova, Italy
- ⁴⁵LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104
- ⁴⁷Istituto Nazionale di Fisica Nucleare Pisa, ^xUniversity of Pisa,
^yUniversity of Siena and ^zScuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- ⁴⁹Purdue University, West Lafayette, Indiana 47907
- ⁵⁰University of Rochester, Rochester, New York 14627
- ⁵¹The Rockefeller University, New York, New York 10021
- ⁵²Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,
^{aa}Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵³Rutgers University, Piscataway, New Jersey 08855
- ⁵⁴Texas A&M University, College Station, Texas 77843
- ⁵⁵Istituto Nazionale di Fisica Nucleare Trieste/Udine,
I-34100 Trieste, ^{bb}University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁶University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁷Tufts University, Medford, Massachusetts 02155
- ⁵⁸Waseda University, Tokyo 169, Japan
- ⁵⁹Wayne State University, Detroit, Michigan 48201
- ⁶⁰University of Wisconsin, Madison, Wisconsin 53706
- ⁶¹Yale University, New Haven, Connecticut 06520
- (Dated: September 4, 2009)

We report on the first direct search for charged Higgs bosons decaying into $c\bar{s}$ in $t\bar{t}$ events produced by $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. The search uses a data sample corresponding to an integrated luminosity of 2.2 fb^{-1} collected by the CDF II detector at Fermilab and looks for a resonance in the invariant mass distribution of two jets in the lepton+jet sample of $t\bar{t}$ candidates. We observe no evidence of charged Higgs bosons in top quark decays. Hence, 95% upper limits on the top quark decay branching ratio are placed at $\mathcal{B}(t \rightarrow H^+b) < 0.1$ to 0.3 for charged Higgs boson masses of 60 to 150 GeV/ c^2 assuming $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1.0$. The upper limits on $\mathcal{B}(t \rightarrow H^+b)$ are also used as model-independent limits on the decay branching ratio of top quarks to generic scalar charged bosons beyond the standard model.

PACS numbers: 11.30.Pb, 12.60.Jv, 14.65.Ha, 14.80.Cp

*Deceased

[†]With visitors from ^aUniversity of Massachusetts Amherst,

The standard model (SM) is remarkably successful in describing the fundamental particles and their interactions. Nevertheless, it is an incomplete theory. An important unresolved question is the mechanism of electroweak symmetry breaking (EWSB). In the SM, a single complex scalar doublet field breaks the symmetry, resulting in massive electroweak gauge bosons and a single observable Higgs boson [1]. To date, the Higgs boson has not been discovered, and consequently the mechanism of EWSB remains in question.

Beyond the SM, many diverse hypotheses with extended Higgs sectors have been proposed to explain EWSB. The simplest extension is a two Higgs-doublet model (2HDM). The minimal supersymmetric standard model (MSSM) employs the type-II 2HDM, where at leading order one doublet couples to the up-type fermions and the other couples to the down-type fermions [2]. The two Higgs doublet fields manifest themselves as two charged Higgs bosons (H^\pm) and three neutral Higgs bosons (h, H, A).

In 2HDM and MSSM, the top quark is allowed to decay into a charged Higgs boson (H^+) [3] and a bottom quark. The tree level branching ratio of top quarks to H^+ , $\mathcal{B}(t \rightarrow H^+ b)$, is a function of the H^+ mass (m_{H^+}) and $\tan\beta$. The parameter $\tan\beta$ is the ratio of vacuum expectation values of the two Higgs doublets. In MSSM, $\mathcal{B}(t \rightarrow H^+ b)$ also depends on extra parameters related to the masses and couplings of the other supersymmetric particles. The $\mathcal{B}(t \rightarrow H^+ b)$ is relatively large if $\tan\beta$ is small ($\lesssim 1$) or large ($\gtrsim 15$) [4]. At low $\tan\beta$, H^+ predominantly decays into $c\bar{s}$ for low m_{H^+} ($\lesssim 130$ GeV/ c^2) and $t^* \bar{b} (\rightarrow W b \bar{b})$ [5] for higher m_{H^+} . In the high $\tan\beta$ region, the H^+ decays into $\tau^+ \nu$ almost 100% of the time.

At Tevatron collider experiments, H^+ searches have been performed for the $H^+ \rightarrow \tau \bar{\nu}$ in $t\bar{t}$ decays. Some searches placed direct upper limits on $\mathcal{B}(t \rightarrow H^+ b)$ by

taking advantage of the expectation that $\mathcal{B}(H^+ \rightarrow \tau \bar{\nu}) = 1.0$ at high $\tan\beta$ [6]. Other searches set limits on the MSSM parameter plane ($m_{H^+}, \tan\beta$) using inclusive H^+ decay branching ratios in the MSSM [7]. The various H^+ final states supplement the SM $t\bar{t}$ decay channels. The previous searches focused on measuring deviations from the SM prediction for the $t\bar{t}$ production and decay, rather than reconstructing H^+ bosons.

In this Letter, we report on the first direct search for $H^+ \rightarrow c\bar{s}$ produced in top quark decays by fully reconstructing the $c\bar{s}$ mass. The final state of $H^+ \rightarrow c\bar{s}$ is mostly two jets, as is the hadronic decay of the W boson [8] in SM top quark decays. The search is performed by looking for a second peak in the dijet mass spectrum (in addition to that from the W boson) in top quark decays.

In the SM, each top quark decays into a W boson and a b -quark exclusively. In this analysis we use the lepton+jets $t\bar{t}$ sample [9], where in the SM one W decays to quarks ($q\bar{q}'$) and the other W decays to $e\bar{\nu}$ or $\mu\bar{\nu}$. Each final-state quark is assumed to form a hadronic jet; the jets are clustered using a cone algorithm with a cone radius $\Delta R (= \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2})$ of 0.4 [10]. This lepton+jets sample has a good signal-to-background ratio for $t\bar{t}$ and is ideal for dijet mass analysis.

The CDF II experiment at the Fermilab Tevatron measures the products of proton-antiproton collisions at $\sqrt{s} = 1.96$ TeV. The lepton momentum is measured using an eight-layer silicon microstrip detector and a cylindrical drift chamber immersed in a 1.4 T magnetic field. The energies of electrons and jets are measured using calorimeters with acceptance up to pseudorapidity as of $|\eta| = 3.6$ [11]. Charged particle detectors outside the calorimeter identify muon candidates up to $|\eta| = 1.0$. Details of CDF II can be found elsewhere [12].

Lepton+jets $t\bar{t}$ events are selected by requiring an electron or a muon with $p_T > 20$ GeV within $|\eta|=1$ and by requiring missing transverse energy larger than 20 GeV to account for the neutrino [13]. Then, the four most energetic jets (called leading jets) within $|\eta| < 2.0$ are required to have $E_T > 20$ GeV after jet energy corrections [10]. In addition, at least two of the leading jets are required to contain a long-lived hadron containing a b -quark [14] by demanding that these jets contain tracks forming a displaced secondary vertex (called a b -tag).

The SM processes are regarded as backgrounds for the H^+ search. The largest background is W bosons in SM $t\bar{t}$ events (92% of the total background). The rest of the SM processes are referred to as non- $t\bar{t}$ backgrounds. These include W +jets, multijets, Z +jets, diboson (WW, WZ, ZZ), and single top events. Details of the non- $t\bar{t}$ background estimation method are given in [14]. Assuming a $t\bar{t}$ cross section of 6.7 pb [15] and a top quark mass of 175 GeV/ c^2 , we expect 152.6 ± 25.0 events from SM $t\bar{t}$ production and 13.9 ± 7.5 events from non- $t\bar{t}$ backgrounds in the 2.2 fb $^{-1}$ CDF II data sample.

The mass of the H^+ candidate is directly reconstructed

Amherst, Massachusetts 01003, ^bUniversiteit Antwerpen, B-2610 Antwerp, Belgium, ^cUniversity of Bristol, Bristol BS8 1TL, United Kingdom, ^dChinese Academy of Sciences, Beijing 100864, China, ^eIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^fUniversity of California Irvine, Irvine, CA 92697, ^gUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^hCornell University, Ithaca, NY 14853, ⁱUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^jUniversity College Dublin, Dublin 4, Ireland, ^kUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^lUniversidad Iberoamericana, Mexico D.F., Mexico, ^mQueen Mary, University of London, London, E1 4NS, England, ⁿUniversity of Manchester, Manchester M13 9PL, England, ^oNagasaki Institute of Applied Science, Nagasaki, Japan, ^pUniversity of Notre Dame, Notre Dame, IN 46556, ^qUniversity de Oviedo, E-33007 Oviedo, Spain, ^rTexas Tech University, Lubbock, TX 79409, ^sIFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain, ^tUniversity of Virginia, Charlottesville, VA 22904, ^uBergische Universität Wuppertal, 42097 Wuppertal, Germany, ^{cc}On leave from J. Stefan Institute, Ljubljana, Slovenia,

using the two jets. The mass resolution is improved by reconstructing the $t\bar{t}$ event as a whole with a kinematic fitter used for the precision top quark mass measurement described in Ref. [12]. The original kinematic fitter is modified for the H^+ search. In the fitter, the lepton, the missing E_T (from a neutrino), and the four leading jets are assigned to the decay particles from the $t\bar{t}$ event, and the quality of the assignment is evaluated using this χ^2 :

$$\chi^2 = \sum_{k=jjb, l\nu b} \frac{(M_k - M_t)^2}{\Gamma_t^2} + \sum_{i=l, 4\text{jets}} \frac{(p_T^{i,\text{fit}} - p_T^{i,\text{meas}})^2}{\sigma_i^2} + \frac{(M_{l\nu} - M_W)^2}{\Gamma_W^2} + \sum_{j=x,y} \frac{(p_j^{UE,\text{fit}} - p_j^{UE,\text{meas}})^2}{\sigma_{UE}^2}. \quad (1)$$

The χ^2 is minimized by constraining leptonic W final state ($l\nu$) to have the W invariant mass (80.4 GeV/ c^2) [16] and both top quark final states ($bl\nu$ and bjj) to have the same top quark mass of 175 GeV/ c^2 . No constraint is imposed on the dijet mass of the hadronic boson (jj). In the mass constraints, the transverse energies of the final-state objects ($p_T^{i,\text{meas}}$) are allowed to vary within measurement uncertainties (σ_i). The unclustered energy ($p_j^{UE,\text{meas}}$) is the sum of measured transverse energies not included in the leading jets E_T and is used to correct the missing E_T . In the jet assignment, b -tagged jets are assigned to the b -quarks. The jets assigned to the b -quarks are called b -jets, and the other two jets are called h -jets. If the $t\bar{t}$ event has more than two b -tagged jets, the jets with the best χ^2 are assigned to b -quarks. Then, we reconstruct the mass of hadronic boson using two h -jets with fit energies ($p_T^{i,\text{fit}}$). In this kinematic event reconstruction, only 55% of the SM $t\bar{t}$ events have correctly matching jets. The wrong jet-parton assignments dominantly come from hard radiation jets which are selected as leading jets and from the falsely b -tagged jets originating from the hadronic decays of W bosons.

The expected dijet mass distributions of H^+ and W in top quark decays are produced using the PYTHIA generator [17] and the full CDF II detector simulation. The ALPGEN generator [18] with the PYTHIA parton shower simulation is used for non- $t\bar{t}$ backgrounds. In the simulation sample, the H^+ is forced to decay solely into $c\bar{s}$ with zero width and with masses ranging from 60 to 150 GeV/ c^2 .

The simulation shows that the reconstructed H^+ has a significant low-mass tail, which is predominantly caused by final-state gluon radiation (FSR) from the hadronic decays of the Higgs boson. The hard FSR results in more than four final-state jets in a lepton+jets $t\bar{t}$ event. To recover the energy loss due to the FSR, the fifth most energetic jet is merged with the closest jet among the four leading ones if the pair has a ΔR distance smaller than 1.0, provided that the fifth most energetic jet has $E_T > 12$ GeV and $|\eta| < 2.4$. Merging the fifth jet results in better jet energy resolution and improves the m_{H^+}

resolution by approximately 5% in more than four final jets events for 120 GeV/ c^2 Higgs sample.

In the CDF II data sample of 2.2 fb $^{-1}$, we observe 200 $t\bar{t}$ candidates in the lepton+jets decay channel. No significant excess is observed in the dijet invariant mass of top quark decays. Figure 1 shows that the observed dijet mass distribution agrees with the SM expectations. Hence, we extract upper limits on $\mathcal{B}(t \rightarrow H^+b)$ using a binned likelihood fit on the dijet mass distribution.




FIG. 1: Observed dijet mass distribution (crosses) compared with background distributions of W bosons (filled) and non- $t\bar{t}$ processes (cross hatched) in CDF II data sample of 2.2 fb $^{-1}$; the background distributions are added on top of each other. An example of the dijet mass distribution from 120 GeV/ c^2 H^+ bosons (bold line) is overlaid assuming $\mathcal{B}(t \rightarrow H^+b) = 0.1$, which is about the 95% C.L. upper limit on $\mathcal{B}(t \rightarrow H^+b)$.

The binned likelihood (LH) function is constructed employing Poisson probabilities:

$$\mathcal{L} = \prod_i \frac{\nu_i^{n_i} \times e^{-\nu_i}}{n_i!} \times G(N_{\text{bkg}}, \sigma_{N_{\text{bkg}}}). \quad (2)$$

The probability of finding events in the mass bin i comes from a set of simulated dijet mass distributions of H^+ , W , and non- $t\bar{t}$ backgrounds. These distributions are called templates. The Poisson probability (P^i) in each bin is computed from the number of observed events, n_i , and from the number of expected events, $\nu_i = P_{H^+}^i \times N_{H^+} + P_W^i \times N_W + P_{\text{bkg}}^i \times N_{\text{bkg}}$, where N_{H^+} , N_W , and N_{bkg} are parameters representing the total number of events in each template category. The minimization of $-\ln \mathcal{L}$ gives the most probable values for N_{H^+} , N_W , and N_{bkg} . In the LH fit, N_{H^+} and N_W are free to vary, however, the non- $t\bar{t}$ background (N_{bkg}) is estimated independently and is allowed to vary within its Gaussian uncertainty ($\sigma_{N_{\text{bkg}}}$). Based on the number of events from the LH fit, a $\mathcal{B}(t \rightarrow H^+b)$ is extracted

assuming $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1$. In Figure 1, dijet mass distributions of the SM events are normalized by the likelihood fit to the observed dijet mass distribution with $\mathcal{B}(t \rightarrow H^+b)$ fixed to 0.

The sources of systematic uncertainty in the extracted $\mathcal{B}(t \rightarrow H^+b)$ include uncertainties in the jet energy scale corrections [10], initial state and final-state radiation, modeling of the non- $t\bar{t}$ background, choice of event generators in simulation. These systematic sources perturb the shape of the dijet mass and cause a shift in the result of the LH fit. The shift in the resulting $\mathcal{B}(t \rightarrow H^+b)$ is estimated using “pseudoexperiments” of the perturbed and unperturbed dijet mass distributions for each systematic source; the pseudoexperiments are generated by the bin-to-bin Poisson fluctuations of the simulated dijet mass distributions. The dominant systematic uncertainty originates from the choice of event generators in the simulation, unless m_{H^+} is close to m_W , in which case the jet energy scale uncertainty dominates. The other systematic uncertainties from data/Monte Carlo differences in b -tagging rates and top quark mass constraints in $t\bar{t}$ reconstruction are negligible compared to the uncertainties from the perturbed dijet mass shape.

The individual systematic uncertainties are combined in quadrature. The total systematic uncertainty ($\Delta\mathcal{B}(t \rightarrow H^+b)$) is represented by a nuisance parameter which adds to the branching ratio and has a Gaussian prior PDF with width $\Delta\mathcal{B}(t \rightarrow H^+b)$. We eliminate this nuisance parameter by Bayesian marginalization [19] and obtain a posterior PDF in $\mathcal{B}(t \rightarrow H^+b)$ assuming a uniform prior PDF in $0 \leq \mathcal{B}(t \rightarrow H^+b) \leq 1$. The expected upper limits on $\mathcal{B}(t \rightarrow H^+b)$ with 95% C.L. are derived from a thousand pseudoexperiments using the SM backgrounds events for each m_{H^+} .

The upper limits on $\mathcal{B}(t \rightarrow H^+b)$ at 95% C.L. show a good agreement between the observation and the SM expectation. The upper limits in Figure 2 includes the systematic uncertainty in $\mathcal{B}(t \rightarrow H^+b)$. Since the LH fit has very little sensitivity for $m_{H^+} \approx m_W$, the upper limits around 80 GeV/c^2 H^+ are omitted in the Figure. The exact values of the upper limits in the Figure 2 are listed in Table I.

$m_{H^+}(\text{GeV}/c^2)$	60	70	90	100	110	120	130	140	150
Expected	0.13	0.19	0.22	0.15	0.13	0.12	0.10	0.10	0.09
Observed	0.09	0.12	0.32	0.21	0.15	0.12	0.08	0.10	0.13

TABLE I: Expected and Observed 95% C.L. upper limits on $\mathcal{B}(t \rightarrow H^+b)$ for H^+ masses of 60 to 150 GeV/c^2 .

This analysis can set model-independent limits for anomalous scalar charged bosons production in top quark decays. Besides the assumption that a scalar boson decays only to $c\bar{s}$ with zero width, no model-specific parameter is used in this analysis. Therefore any generic




FIG. 2: The upper limits on $\mathcal{B}(t \rightarrow H^+b)$ at 95% C.L. for charged Higgs masses of 60 to 150 GeV/c^2 except a region for $m_{H^+} \approx m_W$. The observed limits (points) in 2.2 fb^{-1} CDF II data are compared to the expected limits (solid line) with 68% and 95% uncertainty band.

charged boson would make a secondary peak in the dijet mass spectrum if it decays into a dijet final state like the $H^+ \rightarrow c\bar{s}$ in top quark decays. Here, we extend the search below the W boson mass [20] down to 60 GeV/c^2 for any non-SM scalar charged boson produced in top quark decays, $t \rightarrow X^+(\rightarrow u\bar{d})b$. This process is simulated for the CDF II detector and is similar to $H^+ \rightarrow c\bar{s}$. In the simulation, we obtain a better dijet mass resolution for $u\bar{d}$ decays than for the $c\bar{s}$ decays. The difference in the mass resolution comes from the smaller chance of false b -tagging from light quark final states of X^+ than the $c\bar{s}$ decays, thus resulting in a smaller ambiguity of jet-parton assignments in the $t\bar{t}$ reconstruction. Consequently, the upper limits on $\mathcal{B}(t \rightarrow X^+(\rightarrow u\bar{d})b)$ are lower than the limits on $\mathcal{B}(t \rightarrow H^+(\rightarrow c\bar{s})b)$ regardless of the charged boson mass.

In summary, we have searched for a non-SM scalar charged boson, primarily the charged Higgs boson predicted in the MSSM, in top quark decays using lepton+jets $t\bar{t}$ candidates. This is the first attempt to search for $H^+ \rightarrow c\bar{s}$ using fully reconstructed charged Higgs bosons. In the CDF II data sample of 2.2 fb^{-1} , we find no evidence of charged Higgs bosons in the dijet mass spectrum of the top quark decays. Hence, upper limits on $\mathcal{B}(t \rightarrow H^+b)$ with 95% C.L. are placed at 0.1 to 0.3 assuming of $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1.0$ for charged Higgs masses of 60 to 150 GeV/c^2 . This analysis also yields conservative upper limits on any non-SM scalar charged boson X^+ production from top quarks. Based on simulation, we find that the upper limits on the branching ratio

$\mathcal{B}(X^+ \rightarrow u\bar{d})$ are always better than the upper limits on $\mathcal{B}(H^+ \rightarrow c\bar{s})$.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

-
- [1] P. W. Higgs, Phys. Lett. **12**, 132 (1964); Phys. Rev. Lett. **13**, 508 (1964); Phys. Rev. **145**, 1156 (1966); F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964); G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).
 - [2] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977); J. F. Donoghue and L. F. Li, Phys. Rev. D **19**, 945 (1979).
 - [3] Particles with a superscript (H^+) include the opposite charged particle (H^-), and decay processes such as $t \rightarrow H^+ b$ imply also their charge conjugate processes.
 - [4] M. Carena and H. E. Haber, Prog. Part. Nucl. Phys. **50**, 63 (2003).
 - [5] The t^* is a virtual top quark with an off-shell mass.
 - [6] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. D **62**, 012004 (2000); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 357 (1997); V.M. Abazov *et al.* (D0 Collaboration), arXiv:0906.5326 (2009).
 - [7] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 042003 (2006); B. Abbott *et al.* (DØ Collaboration), Phys. Rev. Lett. **82**, 4975 (1999).
 - [8] The positive and negative charged weak boson (W^\pm) is presented as W in this Letter.
 - [9] In this Letter, the word lepton (l) stands for an electron or a muon. The tau lepton is not referred to as a lepton, but is specified as τ .
 - [10] A. Bhatti *et al.*, Nucl. Instrum. Methods Phys. Res. A **566**, 375 (2006).
 - [11] The detector uses cylindrical coordinates where θ is the polar angle with respect to the proton beam and ϕ is the azimuthal angle. The direction of a particle in the detector is expressed with the pseudorapidity, $\eta = -\ln \tan(\theta/2)$.
 - [12] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **73**, 032003 (2006).
 - [13] Transverse energy is $E_T = E \sin \theta$, where E is the measured energy. Transverse momentum, p_T , of charged particles, is calculated similarly. From energy-momentum conservation, $\sum_i E_T^i \hat{n}_i = 0$ for radial unit vector \hat{n}_i in the azimuthal plane. Missing transverse energy is defined as $-\sum_i E_T^i \hat{n}_i$, representing undetected particles such as neutrinos. The missing E_T is further corrected for the energy and momentum of identified muons.
 - [14] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
 - [15] M. Cacciari *et al.*, J. High Energy Phys. **04**, 068 (2004); N. Kidonakis and R. Vogt, Phys. Rev. D **68**, 114014 (2003).
 - [16] C. Amsler *et al.* (Particle Data Group), Phys. Lett. **B667**, 1 (2008).
 - [17] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. **0605**, 026 (2006). We use PYTHIA version 6.216.
 - [18] M. L. Mangano *et al.*, J. High Energy Phys. **0307**, 001 (2003).
 - [19] Daniel B. Rowe, *Multivariate Bayesian Statistics* (CRC, Chapman & Hall, 2003), p117.
 - [20] LEP Higgs Working Group, hep-ex/0107031v1 (2001). A MSSM charged Higgs mass below m_W is excluded by LEP experiments.